



## Agromet Advisory Bulletin

Date : 11-06-2024

Weather Forecast of **KUHI** Block in **NAGPUR**(Maharashtra) Issued On :2024-06-11(Valid Till 08:30 IST of the next 5 days)

Parameter	2024-06-12	2024-06-13	2024-06-14	2024-06-15	2024-06-16
Rainfall	0.0	0.0	0.0	8.5	2.3
Tmax(°C)	40.6	41.0	39.0	41.5	40.1
Tmin(°C)	29.2	28.0	27.9	27.0	24.8
RH-I(%)	41	46	42	49	60
RH-II(%)	21	19	23	21	24
Wind Speed(kmph)	13	18	21	23	17
Wind Direction(Degree)	281	330	283	290	289
Cloud Cover(Octa)	3	3	3	4	5

### Weather Summary/Alert:

• As per the blocklevel value added forecast given by, IMD, RMC, Nagpur, sky will be partially cloudy during next five days i.e. 12th to 16th, June, 2024. • Very light to light rainfall very likely to occur on 12th, 13th, 14th, 15th and 16th, June, 2024. • Thunderstorm with lightning accompanied with gusty winds (40-50 kmph) likely to occur on 12th and 13th, June, 2024. • Thunderstorm with lightning accompanied with gusty winds (30-40 kmph) likely to occur on 14th, June, 2024. • No large change in maximum and minimum temperature over the Vidarbha during next 5 days.

### General Advisory:

• To prepare kharif vegetable seedlings, sow seeds on raised bed (nursery). Seeds of Chili, Brinjal, and Tomato should be treated with Trichoderma fungicide and seed should be placed at appropriate spacing. • Land preparation should be done (ploughing, harrowing) across the main slop/parallel to counter line to avoid the soil erosion and in situ moisture conservation. • Cows, buffaloes, goats, sheep and other domestic animals should be avoided to graze in the open spaces on days when gustywinds and thunder are predicted. Animals should be kept away from open water sources, rivers or lakes and away from tractors and other metal implements. Sufficient fodder and water should be arranged for the animals in the manger. • Farmers and farm labourers should take care of themselves and livestock keeping in mind the forecast of lightning. Sheltering under trees should be strictly avoided and also livestock should not shelter under trees. • Agricultural produce market committee and private agricultural produce purchase holders should store the agricultural produce brought by the farmers for purchase in the shed instead of in the open. • Marketable vegetables and fruits should be harvested and sold keeping in mind the forecast of gusty winds. Orchards should be supported with bamboo. • Farmers and agricultural labourers should download and use Damini (Lightning Alert) mobile app from Google Play Store and App Store to get early warning of lightning.

### SMS Advisory:

• Farmers and farm labourers should take care of themselves and livestock keeping in mind the forecast of lightning. Sheltering under trees should be strictly avoided and also livestock should not she

### Crop Specific Advisory:

Crop(Varieties)	Crop Specific Advisory
RICE	• Direct Seeded Rice: Direct Seeded Rice should be planted from second week of June to 10th of July.Ploughing twice and then disc harrow or rotavator so that the soil become tilled properly. Short to mid-late rice varieties should be selected.Be careful when sowing, paddy seed should not be sown deeper than 5 cm. • For weed control Pendimethalin 30% E.C. (Pre-emergence) @ 50 ml. Or Pretilachlor 50% E.C.

Crop(Varieties)	Crop Specific Advisory
	(Pre-emergence) @ 20 to 30 ml. in 10 liters of water, spray immediately after sowing if there is moisture and weeding should be done after 30 days. Desi plough, Tifan, Marker, Tractor, Tractor drawn seed drill, Sowing Machine, Drum seeder etc. can be used for paddy sowing. • Green Manuring Crop: If there is abundant water availability in the field, green manure crops should be taken to increase the fertility of the soil, organic matter and the availability of nitrogen and other nutrients. • Green manure to be planted in the field: - Green manure is sown in the field continuously or as an intercrop in fruit crops and in the same field these crops are mixed in the field before flowering. These types of green manure include hemp, cluster bean, cowpea, dhaincha, green gram, matki, black gram, etc. • Kharif Paddy: • Seed Germination test should be done if you want to use own seed. Use those seed for sowing if found germination more than 70 to 80% and use seed accordingly. Rice varieties recommended for East Vidarbha should be used for cultivation. Seed treatment should be done before sowing. For seed treatment, dissolve 300 g of salt (3%) in 10 liters of water and pour seeds in it. Once the solution is stable, remove the floating diseased seeds through a sieve and burn them. Wash the healthy seeds at the bottom 2-3 times with clean water and dry them in the shade for 24 hours. Before sowing, seeds should be treated with fungicide and Trichoderma bio-fungicide should be applied by rubbing 4 to 5 gm per kg of seed.
COTTON	• Avoid pre-monsoon sowing of cotton crop. • Crop rotation to be followed in the fields that were heavily infested with pink bollworm during last season to break the life cycle of pink bollworm. • Cotton is the only host of pink bollworm, therefore crop rotation helps to break the life cycle of this pest. • Crop rotation is very effective in checking the infection of soil borne diseases and nematodes in disease prone fields.
SOYABEAN	• As per suitability, farmers are advised to run the sub-soiler machine once in five years, at an interval of 10 m which facilitates the breaking of hard soil pan thereby increasing the rainwater infiltration and helps in case of drought. • Farmers are advised to apply deep summer ploughing once in 3 years. After this, the field should be prepared using criss-cross harrowing followed by planking. • Apply well-decomposed FYM @ 5- 10 t/ha or Poultry Manure @ 2.5 t/ha before the last harrowing. • Farmers are advised to follow Broad Bed Furrow (BBF) or Ridge & Furrow methods for sowing. • Ensure availability of other critical inputs like fertilizers, weedicides, fungicides, and cultures for seed treatment etc. • Ensure the seed quality (minimum 70% germination) by carrying out germination test of the available soybean seed.
GROUNDNUT	• For kharif groundnut cultivation, medium, well drained, humus rich, sand mixed clay soil and organic matter rich soil should be select. For groundnut cultivation, soil layer of 15 to 20 cm should be humorous. Apply 5 tons of well decomposed manure before the last harrowing.
PIGEON PEA (RED GRAM/ARHAR)	• Agricultural land should be prepared for cultivation of Arhar crop. Good quality seeds of Arhar crop should be procured from authorized seller/certified seed holder only. Rhizobium or Phosphorus Dissolving Fertilizer (PSB) {250 gm per 10 kg of seed) should be used for seed treatment. Recommended varieties of Arhar are- AKT-8811, PKV-TARA, BSMR-853, BSMR-736 and ICPL-87119 (Aasha).
GREEN GRAM	• PKV Moog-8802, PKV Greengold (AKM 9911) varieties of green gram should be select for cultivation of green gram crop,
BLACK GRAM	• For cultivation of Black gram crop, TAU-1, TAU-2, PKV udid-15 and PKV black gold varieties should be select,

#### Horticulture Specific Advisory:

Horticulture(Varieties)	Horticulture Specific Advisory
MANDARIN ORANGE	• For establishment of new orchard suitability of the soil needs to be examined. The soil depth should be a minimum of 1 meter but equally high-performance orchards can be established on soil depth measuring even less than 1 meter provided precision based cultural practices including the fertilization programme are adopted. The soil site should preferably be well drained, clay less than 60%, pH less than 8.3, free calcium carbonate less than 12 % and water table of the area more than 2 meters should be selected. • The pits excavated during the month of May may be filled up by a mixture containing 2 parts soil, one-part sand and 1 parts farmyard manure. In addition, add 1 kg Single super phosphate, 1 kg of neem cake and 100 g Chlorpyrifos powder in each pit. • Nurseryman of citrus should start filling polythene bags with the mixture of garden soil, FYM and sand. Proper perforation for water should be

Horticulture(Varieties)	Horticulture Specific Advisory
	done on the bags adequately. • Drainage channels (trenching size: 30cm depth, 45 cm top width and 30 cm bottom width) after each 2 rows of plants along the slope should be prepared to remove excess water from the orchard in rainy season. • For effective management of gummosis of Nagpur mandarin gummosis-affected portion from where gum is oozing needs to be scraped out with sharp knife, taking care not to damage the wood. Cleaned portion pasting with Metalaxyl MZ* (50 g/l) or Fosetyl-A1* (50 g/l) at initiation of disease and second after 30 days from initiation of the disease. • For controlling citrus mealy bugs, the basin around the tree trunk should be earthed up. Wrap sticky band by smearing with mobile oil around the tree trunk and destruction of ant's nests should be done and spraying of Chlorpyrifos 20 EC @ 5ml/liters of water and apply in ant nests. For control of mealy bug Chlorpyrifos 20 EC @ 2ml/liters of water or dimethoate 2ml/litre of water should be done on the tree leaves and trunk.
CABBAGE	• Carry out harvesting of market ready curd and head of cabbage and cauliflower respectively. • Considering the prevailing cloudy weather, for management of aphids on Cabbage & Cauliflower, spray Dimethoate 30 EC 13 ml or Cyantraniliprole 10.26 OD 12 ml per 10 liters of water. • Considering the prevailing cloudy weather, for management of Diamond Back Moth on Cabbage & Cauliflower, Spraying should be done of Chlorpyrifos 20 EC 40 ml or Spinosad 2.5 SC 12 ml or Fenvalerate 20 EC 5 ml per 10 liters of water. • Spraying operation should be done during the clear and calm weather.
TOMATO	• Considering the gusty winds and rainfall forecast, carry out harvesting of market ready fruits and support the plant with bamboo. • In tomato, if leaf minor, aphids, thrips, white fly and fruit borer crosses ETL spraying should be done of Cyantraniliprole 10.26 OD at the rate of 18.0 ml per 10 litres of water during clear weather condition.

#### Live Stock Specific Advisory:

Live Stock(Varieties)	Live Stock Specific Advisory
BUFFALO	• Offer clean and cold water (stored in earthen pot) to the animals. Those who have cultivated berseem, should be done top dressing of 10 kg N/ha after each cutting to encourage quick growth and high yield. Deworming of livestock should be done by seeking advice of Veterinary practitioner.
COW	• Dairy cows should be given 8 kg dry fodder and 6 kg green fodder and 1 kg feed. Maize, sorghum, millet, wheat bran should be used in mixing in preparation diet. Add 20 gms per kg of mineral mixture in the diet.