



Gramin Krishi Mausam Sewa
Agromet Advisory Bulletin for NAGPUR District
 Central Institute for Cotton Research,
 Nagpur



Agromet Advisory Bulletin

Date : 2024-10-29

Weather Forecast of District NAGPUR (Maharashtra) Issued On : 2024-10-29 (Valid Till 08:30 IST of the next 5 days)

Parameter	2024-10-30	2024-10-31	2024-11-01	2024-11-02	2024-11-03
Rainfall(mm)	12.3	0.0	0.0	0.0	0.0
Tmax(°C)	34.8	34.7	34.6	34.7	34.8
Tmin(°C)	21.6	21.8	21.4	20.2	19.1
RH-I(%)	89	90	96	96	96
RH-II(%)	58	62	78	78	78
Wind Speed(kmph)	4	3	4	5	5
Wind Direction(Degree)	53	137	46	220	220
Cloud Cover(Octa)	5	5	5	4	3

Weather Summary/Alert:

• As per the district level value added forecast given by, IMD, RMC, Nagpur, sky will be clear to partly cloudy during next five days i.e. to 30th, October to 03rd, November, 2024. • Very light to light rainfall very likely to occur at isolated places on 30th, October, 2024. • Weather is very likely to be dry on 31st, October, 01st, 02nd and 03rd, November, 2024. • Thunderstorm with lightning is likely to occur at isolated places on 30th, October, 2024.

General Advisory:

• Considering the very light to light rainfall forecast at isolated during 24 hours and dry weather for next 4 days, it is advised to give the priority for the harvesting and threshing of mature crop and continue the sowing of Rabbi Crop, intercultural operations, agrochemicals application during next 5 days. • Cows, buffaloes, goats, sheep and other domestic animals should be avoided to graze in the open spaces on days when gusty winds and thunder are predicted. Animals should be kept away from open water sources, rivers or lakes and away from tractors and other metal implements. Sufficient fodder and water should be arranged for the animals in the manger. • Farmers and farm laborers should take care of themselves and livestock keeping in mind the forecast of lightning. Sheltering under trees should be strictly avoided and also livestock should not shelter under trees. • Agricultural produce market committee and private agricultural produce purchase holders should store the agricultural produce brought by the farmers for purchase in the shed instead of in the open. • Marketable vegetables and fruits should be harvested and sold keeping in mind the forecast. • Dry land wheat crop should be sown in the second fortnight of October. • Sowing of irrigated linseed should be done up to 07th, November. • Sowing of safflower under irrigation can be done till the end of October. • Before sowing of gram seed, seed treatment should be done of 5 gm of Trichoderma or 2 gm of Thirum + 2 gm of Carbendazim per kg of seed, followed by 250 gm of Rhizobium (Nitrogen Fixing Bacteria) and 250 gm of P. S. B. (Phosphorus Solubilizing Bacteria) seed treatment should be done by mixing cold solution of jaggery per 10 kg seed. After seed treatment the seeds should be dried in the shade for an hour and then sown. • Seed treatment should be given to chickpea seed prior to sowing to avoid fungal diseases, for this purpose recommended fungicide is Tebuconazole 5.4 % w/w FS @ 0.4ml/kg

SMS Advisory:

• Agricultural produce market committee and private agricultural produce purchase holders should store the agricultural produce brought by the farmers for purchase in the shed instead of in the open.

Crop Specific Advisory:

Crop (Stage)	Crop Specific Advisory
BENGAL GRAM/ CHICK PEA	<p>• Deshi varieties and seed rate of chickpea for sowing- Hirawa Chafa (AKGS-1), Vijay and ICCV-10 (50-60 kg/ha seed), PKV Harita (AKG- 9303-12) and JAKI- 9218 (75-85 kg/ha seed), Kabuli varieties and seed rate of chickpea, PKV Kabuli-2 & PKV Kabuli-4 (110-115 kg/ha seed), Pink chickpea variety and seed rate, Gulak-1 (75-85 kg/ha seed) are recommended for sowing up to second fortnight of October to 15th November and sowing of PKV Kanchan (AKG- 1909) (50-60 kg/ha seed) variety of chickpea up to second fortnight of October to 15th November. • Before sowing of gram seed, seed treatment should be done of 5 gm of Trichoderma or 2 gm of Thirum + 2 gm of Carbendazim per kg of seed, followed by 250 gm of Rhizobium (Nitrogen Fixing Bacteria) and 250 gm of P. S. B. (Phosphorus Solubilizing Bacteria) seed treatment should be done by mixing cold solution of jaggery per 10 kg seed. After seed treatment the seeds should be dried in the shade for an hour and then sown. • Seed treatment should be given to chickpea seed prior to sowing to avoid fungal diseases, for this purpose recommended fungicide is Tebuconazole 5.4 % w/w FS @ 0.4ml/kg to manage Root rot and Wilt disease.</p>
WHEAT	<p>• Medium heavy, deep and well-drained soil should be selected for wheat crop. When preparing the land for wheat crop, 15 to 20 cm deep ploughing should be done. The soil should be harrowing by giving 2 to 3 shifts. Clean the field by removing the previous crop debris and stick waste. Land should be as level as possible so that further irrigation can be managed. Dryland wheat crop should be sown in the second fortnight of October. Use 75 kg seed per hectare for sowing of dryland wheat. Dryland wheat should be sown with sufficient moisture in the soil and adequate care should be taken to ensure that the sown seeds get sufficient soil moisture contact. For sowing of dryland wheat, spacing between two rows should be 23 cm. Care should be taken not to fall deeper than 5 to 6 cm at the time of sowing wheat. • Varieties should be AKDW 2997-16 (Sharad), PDKV Washim (WSM-1472), MACS 1967 and NI 5439 for dry land wheat sowing. PDKV Washim (WSM-1472) should be sown under limited availability of irrigations. Varieties should be AKDW 1071 (Purna), AKDW 3722 (Vimal), HD 2189 and HD 2380 for timely sowing of irrigated wheat, Varieties should be PDKV Sardar (AKAW 4210-6), AKAW 4627, AKAW-381, AKAW 1071 (Purna) and HI 977 for late sowing of irrigated wheat. • Before sowing, wheat seeds should be treated Azotobacter (Nitrogen fixing bacteria) and Phosphorus solubilizing bacteria fertilizer at the rate of 250 g / 10 to 12 kg of seed. • Seed treatment should be given to wheat seed prior to sowing avoid fungal diseases and pest incidence, for this purpose recommended fungicide and insecticides are Carboxin 75 % WP @ 2-2.5 gm/kg of seed to manage Bunt, Flag smut and Loose smut disease or Difenoconazole 3 % WS @ 2 gm/kg to manage Loose smut disease or Tebuconazole 5.4 % w/w FS @ 0.3 ml/kg of seed to manage Loose smut and Flag smut disease or Carboxin 37.5% + Thiram 37.5% WS @ 3 gm/kg of seed to manage Loose smut disease or Imidacloprid 18.5 % + Hexaconazole 1.5% FS @ 2 ml/kg of seed to manage the Rust and Smut diseases as well as Termite and Aphid incidence.</p>
SAFFLOWER	<p>• Sowing of safflower under irrigation can be done till the end of October. • If the seeds are soaked overnight and sown, germination is quick and good. • While sowing, the distance between two rows should be 45 cm. • Safflower crop can be taken as an intercrop with chickpea, linseed system in the ratio of Safflower + Chickpea (6:3) or Safflower + Linseed (3:3). • 40 kg of Nitrogen (194 kg of ammonium sulphate or 87 kg of urea) + 25 kg of Phosphorus (156</p>

Crop (Stage)	Crop Specific Advisory
	kg of single super phosphate) per hectare should be applied to dryland Safflower crop.
SOYABEAN	<ul style="list-style-type: none"> • The harvested crop must be threshed after sun drying. If the threshing is not done immediately, it should be stored at safe place protecting from rains. • If the produce is to be used for seed purpose in the next season, farmers are advised to thresh the soybean at 350 to 400 RPM thresher to avoid the loss of seed germination. • When storing soybean seeds, the moisture content should not exceed 8%. A 100 kg bag should not be added more than 5 times. Do not hit the seed bag.
COTTON	<ul style="list-style-type: none"> • If the incidence of internal boll rot/bacterial boll rot is noticed in cotton at squaring, flowering and boll development stage due to high humidity, cloudy weather and continuous rainfall during previous week, for management it is advised to collect and destroy the dried petals sticking to the developing bolls should be removed. Avoid indiscriminate use of nitrogenous fertilizers. Restrict excess vegetative growth of the cotton crop. Facilitate proper drainage in the field to avoid water logging in the field. A prophylactic spray of Copper Oxychloride 50 WP @ 25g/10 L is suggested during early boll developmental stages at 15 days interval. Manage sucking pests with spray of recommended insecticides. • If the incidence Target leaf spot, Alternaria leaf spot, Myrothecium leaf spot, External fungal boll rot was noticed in cotton crop due to due to high humidity, cloudy weather and continuous rainfall during previous week, it is advised to take the prophylactic spray of Propineb 70 WP @25-30 g Or Azoxystrobin 18.2 % w/w + Difenconazole 11.4 % w/w SC @ 10 ml Or Fluxapyroxad 167 g/l + Pyraclostrobin333 g/l SC@ 6g Or Carbendazim 50 WP @ 20 gm Or Propiconazole 25 EC @10 ml Or Pyraclostrobin 5% + Metiram 55% WG @ 20 g per 10 litres water. • If the incidence of adult white fly is noticed, on crossing ETL spray Diafenthuron 50% WP @ 12g/10L (600 g/ha) Or Afidopyropen 50 g/L @ 20ml/10L (1000 ml/ha) Or Dinotefuran 20 SG @ 3g/10L (150g/ha) Or Flonicamid 50 WG @4g/10L (200 g/ha) or Clothianidin 50%WDG 1ml/10L (50ml/ha) by judging the local calm and clear weather condition. • At boll formation stage, farmers are advised to inspect the presence and damage of pink bollworm by plucking 20 green bolls from different plants randomly (one boll per plant). If ETL crossed i.e. >10% damaged flowers (Rosette flowers) or 10% damaged green bolls (at least two out of 20 bolls having white or pink larvae or exit holes) and or 8 moths catch per pheromone trap for consecutive 3 days, spray Profenofos 50 EC @ 30 ml/10L (1500 ml/ha) Or Emamectin benzoate 5 SG @ 5g/10L (250 g/ha) Or Indoxacarb 14.5 SC @10ml/10L (500ml/ha) Or Chlorpyrifos 20 % EC @ 25ml/10L (1250 ml/ha)by judging the local calm and clear weather condition.
RICE	<ul style="list-style-type: none"> • Keeping view in weather, Farmers are advised to harvest of mature rice crop should be done close to the ground so that the pupation of stem borer is eliminated and the incidence of this pest in summer paddy crop is reduced. • After harvesting, crop should be dried in the field for 2-3 days and thereafter threshing should be done. • False smut infected panicles should be removed separately, buried in a deep pit or burn it so that the disease does not spread elsewhere. • Leaf roller/folder: - Nitrogenous fertilizers should be used in a balanced manner. Infected leaves wrapped by larvae should be collected and destroyed with larvae. Beauveria bassiana as a bio-insecticide 1.15 % @ 2.25 kg/ha or Azadirachtin 0.15 % @ 30-50 ml. or Quinalphos 20% F @25 ml. or Indoxacarb 15.8 % @ 4.0 ml. spray by mixing in 10 liters of water. • Plant hoppers: - Although rice crop is prone to plant hoppers, use Metarhizium anisopliae as a bio-insecticide 1.15% @ 2.5 kg/ha. Buprofezin 25% @16 ml for control as soon as the level of financial loss is exceeded. or Imidacloprid 17.8 SL.@ 2.0 ml. or Fipronil 5 SC@ 20 ml. or Flonicamid 50 WG @ 3.0 gm Mix in 10 liters of water and spray. • Disease management: - • Blast and Neck blast: - Spray Hexaconazole 5% EC @ 20 ml. or Mencozeb 75% @ 30 gm per 10 liters of water. • Bacterial leaf blight: - Spray Copper hydroxide 53.8% DF @ 30 gm + Streptocycline 1.5 gm per 10 liters of water. • False smut: - At

Crop (Stage)	Crop Specific Advisory
	50% flowering stage of paddy crop spray Copper hydroxide 77% WP @ 30 gm per 10 liters of water in the afternoon.

Horticulture Specific Advisory:

Horticulture (Stage)	Horticulture Specific Advisory
LINSEED	<ul style="list-style-type: none"> Sowing of irrigated linseed should be done up to 07th, November. Before sowing, seed treatment should be done of Carbendazim 2 gm or Thirum 3 gm per kg of seed. After 3 hours, seed treatment should be done at the rate of 20 g of Azotobacter and 20 g of phosphorus solubilizing bacteria per kg of seed.
MANDARIN ORANGE	<ul style="list-style-type: none"> Fruit fly- To attract fruit fly males, fruit fly traps (methyl eugenol) at the rate of 25 per hectare should be hung on trees in the orchard about 2 months before harvesting. The fallen fruits in the orchard should be picked and destroyed and the orchards should be kept clean. The pupal stage of the fruit fly is 2 to 3 cm deep in the soil. The soil under the tree should be moved or hand weeding. Fruit Sucking Moth- To manage Fruit Sucking Moth it is advised to destroy host weed other than Citrus crop e.g. Gulvel, Vasanvel, Chandvel etc. The larval stage of this pest lives on the host plants. Generally, in the evening time (7 to 11 pm) the grass should be burnt and smoked on the orchard's embankment. At the time of fruit ripening, a mercury lamp should be placed in the four corners of the garden as well as in the center and kerosene should be poured in a wide vessel under the lamp. Prepare poisonous baits for fruit sucking moths and keep them in the garden. For this Malathion 50 EC 20 ml + 200 gm jaggery + fallen fruit juice (400 to 500 ml) mixed with 2 liters of water and put two baits each in two wide mouth bottles and keep one in every 25 to 30 plants. When the fruit turns from green to yellowish colour, spray with Nimboli oil (neem oil) or mineral oil mixed with 10 ml per liter of water at an interval of 10-15 days until the fruit is harvested. The fallen fruits in the orchards should be picked and buried in gravel to keep the garden clean. (Note: • Label claim is not recommended; based on research, # Central Pesticides Board, New Delhi recommended) (Source: AICRP on Fruits, Dr. PDKV, Akola) Colletotrichum stem end rot or "Deth Sukhi"- For fruit rot caused by Colletotrichum stem end rot, spray Bordeaux mixture 0.6 per cent or copper oxychloride 50 WP * 2.5 g per liter or Azoxystrobin + Difenconazole 1 ml per liter of water. Greasy Spot- For faster decomposition of fallen leaf litter, other beneficial fungi (bio-decomposer mixed with cow dung (1 kg / tree)) should be used. Zineb * 68 % (20 g/10 liter of water) or horticultural mineral oil @ 2 % 200 ml / 10 liters of water) or pre-mixed fungicide Hexaconazole 4% + Zineb 68 % WP @ 15 gm / 10 liters of water should be sprayed. Application of mineral oil or fungicides reduces spore penetration into the leaf and also reduces spore germination. If the fungus is infected, the development of symptoms is prevented or delayed as well as the severity of the oily spot. Brown rot (brown rot on fruit) - First of all, fallen leaves and fruits should be disposed of on the trees and not allowed to remain in the field; otherwise the disease will increase in severity and spread rapidly. Keep beds clean. Do not place piles of fruit anywhere in the orchard as they serve to spread disease. As a preventative measure to prevent leaf drop and fruit rot caused by Phytophthora fungus, whole plant should be sprayed with Fosetyl AL* 2.5 gm or copper oxychloride * 50 WP 3 gm per liter of water. While sprayi
CHILLI	<ul style="list-style-type: none"> Due to alternative low and high temperature coupled with high humidity, if the powdery mildew disease is noticed on chilli crop, spraying should be done of any of following fungicide, Hexaconazole 75 % WG @ 66.7 gram or Tebuconazole 25% WG @ 500-750 gram or Azoxystrobin 8.3 % + Mancozeb 66.7 % WG @ 1500 gram or Boscalid 25.2% + Pyraclostrobin 12.8 % WG @ 600 gram or Carbendazim 12 % + Mancozeb 63 % WP @ 750 gram or Kresoxim-Methyl 15 % + Chlorothalonil 56 % WG @ 1000 gram or Tebuconazole 10 % WP + Sulphur 65 % WG @ 1250 gram or Tebuconazole

Horticulture (Stage)	Horticulture Specific Advisory
	50 % + Trifloxystrobin 25% WG @ 250 gram mixed with in 500 litre of water per acre for management of disease.

Live Stock Specific Advisory:

Live Stock	Live Stock Specific Advisory
COW	<ul style="list-style-type: none"> • Lucerne and Berseem fodder crops should be planted for availability of nutritious fodder for animals. Keep animal shed clean, dry and well ventilated. Deworming should be done, if not done earlier within 3 months. Disinfection/fumigation of shed should be done by using formalin. Include dry fodder/feed in the diet of small and large ruminants to avoid diarrhoea/indigestion. Provide complete ration to the livestock. Apply recommended doses of fertilizers to fodder crop.
GOAT	<ul style="list-style-type: none"> • Vaccinate the goat against FMD, HS, PPR and enterotoxaemia. Offer clean and cold water (stored in earthen pot) to the animals.

Others (Soil / Land Preparation) Specific Advisory:

Others (Soil / Land Preparation)	Others (Soil / Land Preparation) Specific Advisory
GENERAL ADVICE	<ul style="list-style-type: none"> • Garlic, Radish, Fenugreek, Spinach, Shravan Ghewda, Potato, Carrot, Guar, Pea etc. should be cultivated.